arXiv:cond-mat/9501032v2 13 Feb 1995

IPT-EPFL preprint

A Remark On Interacting Anyons In Magnetic Field

D. F. Wang and C. Gruber

Institut de Physique Théorique

École Polytechnique Fédérale de Lausanne

PHB-Ecublens, CH-1015 Lausanne-Switzerland.

In this remark, we note that the anyons, interacting with each other through pairwise potential $\sum_{i\neq j} V(|\vec{r}_i - \vec{r}_j|)$ in external magnetic field, exhibits a quantum group symmetry $U_q(sl(2))$.

PACS number: 71.30.+h, 05.30.-d, 74.65+n, 75.10.Jm

Typeset Using *REVTEX*

There have been considerable interests in the quantum mechanics of anyons in two dimensions. Anyons have been suggested to be relevant to various interesting systems, such as the quasiparticles in the Fractional Quantum Hall fluids, the solitons in the nonlinear sigma model [1–10]. Due to the nontrivial statistical interaction between the anyons, elementary quantum mechanics of free anyon gas is far from being understood fully [2,3]. For free anyons, the system is found to exhibit the conformal symmetry [5]. Attempts have been made to find the energy spectrum of the free anyon gas, however, only in the case of two anyons, the complete solution is available by far [2,3]. In presence of external magnetic field, the ladder operator approach can provide us with some portion of the energy spectrum and the wavefunctions of the anyon gas [9]. In this letter, we present a simple quantum group symmetry for interacting anyons in external magnetic field.

The anyons in external magnetic field are described in terms of the following Hamiltonian:

$$H = \sum_{i=1}^{N} \frac{1}{2} [\vec{p}_i - \alpha \sum_{j(\neq i)=1}^{N} \frac{\vec{k} \times \vec{r}_{ij}}{|\vec{r}_{ij}|^2} - \frac{eB}{2} \vec{k} \times \vec{r}_i]^2, \tag{1}$$

where we assume that N anyons move in a two dimensional infinite plane, and the statistical parameter α is within the range $[-1, 1]$. For the constant external magnetic field $\vec{B} = B\vec{k}$, we choose the symmetric gauge, such that the vector field is $\vec{A}_i = \frac{B}{2}\vec{k} \times \vec{r}_i$. For these anyons, the bosonic wavefunctions are used as their base functions,

$$\phi(\cdots, \vec{r}_i, \cdots, \vec{r}_j, \cdots) = \phi(\cdots, \vec{r}_j, \cdots, \vec{r}_i, \cdots), \tag{2}$$

and the anyons are composite particles of the bosons attached with flux. The anyon spectrum is determined by the following eigenequation:

$$H\phi(\vec{r}_1, \vec{r}_2, \cdots, \vec{r}_N) = E\phi(\vec{r}_1, \vec{r}_2, \cdots, \vec{r}_N). \tag{3}$$

For the anyons, one may perform the following well-known singular statistical gauge transformation $U = e^{-i\alpha \sum_{k<l} \theta_{kl}}$:

$$\phi'(\vec{r}_1, \vec{r}_2, \cdots) = e^{i\alpha \sum_{k<l} \theta_{kl}} \phi(\vec{r}_1, \vec{r}_2, \cdots), \tag{4}$$

where θ_{ij} is the angle between the vector $\vec{r}_{ij}$ and the x-axis, $\vec{r}_{ij} = \vec{r}_i - \vec{r}_j = r_{ij}(\cos\theta_{ij}\vec{x} + \sin\theta_{ij}\vec{y})$. The eigenenergy equation can be translated into

$$H'\phi' = E\phi', \tag{5}$$

with

$$H' = U^{\dagger}HU = \sum_{i=1}^{N}\frac{1}{2}(\vec{p}_i - \frac{eB}{2}\vec{k}\times\vec{r}_i)^2. \tag{6}$$

For the multivalued wavefunction ϕ', additional phase factor shows up when one interchanges two patricles,

$$\phi'(\cdots, x_i, \cdots, x_j, \cdots) = e^{\pm i\alpha\pi}\phi'(\cdots, x_j, \cdots, x_i, \cdots). \tag{7}$$

One can see that the wavefunction ϕ' is invariant under the change $\alpha \to \alpha + 2$. In particular, when the statistical parameter $\alpha = 1$, the wavefunction ϕ' is antisymmetric, describing N fermions in external magnetic field. For N fermions, the energy spectrum of the system is given by the sum of single particle energies, as long as the Pauli exclusion principle is properly taken into account when one constructs the Slater determinant. For one electron in external magnetic field, we have the well-known Landau problem in two dimenions [17], which is also the problem of Integer Quantum Hall Effect when the magentic field B is so strong, compared to the electron-electron Coulomb interaction and the disorder in real two dimensional electron gas, with small ratio of electron-electron interaction to disorder.

In the Landau problem, the system is not translational invariant. However, it has been known for a long time that the system is invariant under the magnetic translation group. Considering the Hofstadter problem (a free electron hopping on two dimensional lattice in constant magnetic field), Wiegmann and Zabrodin show that the magnetic translations can be constructed with the generators of the quantum group $U_q(sl(2))$ [12]. Applying the representation, they obtain the Bethe-Ansatz. Quite recently, similar discussions have been carried out for the Landau problem, where the quantum group symmetry $U_q(sl(2))$, the W algebra and the area-presevering differomorphism were studied [11].

When one has arbitrary statistical parameter $-1 < \alpha < 1$, the system consists of particles of intermediate statistics. The anyons are interacting with each other through nontrivial statistical gauge term, and the eigenenergy problem can no longer be reduced to a single particle problem. However, in the following, we would like to demonstrate that the system of the anyons still exhibits a magnetic translation group, from which the quantum group symmetry $U_q(sl(2))$ can be constructed. In fact, one will see that the quantum group symmetry will survive, even when the anyons are interacting with each other through pairwise potential $\sum_{i\neq j} V(r_{ij})$.

In presence of pairwise interaction between the anyons, the Hamiltonian for the anyons takes the following form:

$$H_N = \sum_{i=1}^{N} \frac{1}{2}[\vec{p}_i - \alpha \sum_{j(\neq i)=1}^{N} \frac{\vec{k}\times\vec{r}_{ij}}{|\vec{r}_{ij}|^2} - \frac{eB}{2}\vec{k}\times\vec{r}_i]^2 + \sum_{i\neq j} V(r_{ij}). \tag{8}$$

With the singular statistical gauge transformation U, the Hamiltonian becomes

$$H'_N = U^\dagger H_N U = \sum_{i=1}^{N} \frac{1}{2}(\vec{p}_i - \frac{eB}{2}\vec{k}\times\vec{r}_i)^2 + \sum_{i\neq j} V(r_{ij}). \tag{9}$$

Let us construct following generators of magnetic translation:

$$T_{\vec{\zeta}} = \prod_{k=1}^{N} \exp[\zeta_1(\partial_{x_k} + eBiy_k/2) + \zeta_2(\partial_{y_k} - eBix_k/2)], \tag{10}$$

where $\vec{\zeta} = \zeta_1\vec{x} + \zeta_2\vec{y}$ is a two dimensional vector, and the symmetric gauge for the vector field is used, $\vec{A}_i = \frac{B}{2}\vec{k}\times\vec{r}_i$. The generator $T_{\vec{\zeta}}$ is the product of the generators of magnetic translation of each single particle. They satisfy the following group property [16]:

$$T_{\vec{\zeta}} T_{\vec{\eta}} = \exp[-iN\frac{\vec{B}}{2}\cdot(\vec{\zeta}\times\vec{\eta})]\ T_{\vec{\zeta}+\vec{\eta}}. \tag{11}$$

The generators commute with the pairwise potential term,

$$[T_{\vec{\zeta}}, \sum_{i\neq j} V(|\vec{r}_{ij}|)] = 0. \tag{12}$$

One can verify these generators commute with the Hamiltonian H'_N given by Eq. (9),

$$[H'_N, T_{\vec{\zeta}}] = 0. \tag{13}$$

Moreover, with the simple relation $[\sum_{i=1}^{N}\partial_{x_i}, \sum_{k<l}\theta_{kl}] = 0$ and $[\sum_{i=1}^{N}\partial_{y_i}, \sum_{k<l}\theta_{kl}] = 0$, one can check that the singular statistical gauge transformation U commutes with the generators of magnetic translation group,

$$[T_{\vec{\zeta}}, U] = 0. \tag{14}$$

With these equations, we therefore conclude that the magnetic translation group generators also commute with the original anyon Hamiltonian Eq.(8)

$$[H_N, T_{\vec{\zeta}}] = 0, \tag{15}$$

i.e., the anyon gas is invariant under the magnetic translation.

With the generators of magnetic translation group, one can construct the generators of the quantum group $U_q(sl(2))$ as follows [12]:

$$\begin{aligned}
J_+ &= \frac{1}{(q-q^{-1})}(\alpha T_{\vec{a}} + \beta T_{\vec{b}}), \\
J_- &= \frac{1}{(q-q^{-1})}(\gamma T_{-\vec{a}} + \delta T_{-\vec{b}}), \\
q^{2J_3} &= T_{\vec{b}-\vec{a}}, \\
q^{-2J_3} &= T_{\vec{a}-\vec{b}},
\end{aligned} \tag{16}$$

where $q = \exp[iN\frac{\vec{B}}{2}\cdot(\vec{a}\times\vec{b})]$, $\alpha\delta = \beta\gamma = -1$, and $\vec{a}, \vec{b}$ are two vectors defined in the two dimensional plane. Using the group property of the magnetic translation generators, one can check that the following commutation relations of the $U_q(sl(2))$ [13–15] quantum group hold

$$\begin{aligned}
q^{J_3} J_\pm q^{-J_3} &= q^{\pm 1} J_\pm \\
[J_+, J_-] &= \frac{q^{2J_3} - q^{-2J_3}}{q - q^{-1}}.
\end{aligned} \tag{17}$$

The generators for the quantum group $U_q(sl(2))$ also commute with the original anyon Hamiltonian Eq. (8), and we therefore conclude that the interacting anyons in magnetic field also exhibits a quantum group $U_q(sl(2))$,

$$
\begin{aligned}
&[H_N, J_\pm] = 0, \\
&[H_N, J_3] = 0. \qquad (18)
\end{aligned}
$$

The above two equations are the main results we would like to make remarks of in this letter. Finally, we note that the quantum group symmetry is also valid for the interacting electron system in strong magnetic field. The many particle system is the problem of Fractional Quantum Hall when the ratio of disorder to the electron-electron interaction is small. For the interacting electron system, the Hamiltonian is given by

$$
H = \sum_{i=1}^{N} \frac{1}{2}(\vec{p}_i - e\frac{\vec{B}}{2} \times \vec{r}_i)^2 + \sum_{i \neq j} \frac{e^2}{r_{ij}}, \qquad (19)
$$

and the wavefunction ϕ' is totally antisymmetric when interchanging any pair of particles, which corresponds to $\alpha = 1$.

In summary, for arbitrary intermediate statistics, the interacting anyon gas in external magnetic field is invariant under one magnetic translation group, as the singular statistical gauge transformation commutes with such magnetic translations. The interacting anyons hence exhibits a quantum group symmetry $U_q(sl(2))$, which can be readily constructed from the magnetic translation group.

We would like to thank the Swiss National Science Foundation for the financial support. We also would like to thank Ian I. Kogan for communication.

REFERENCES

[1] D. Arovas, J. R. Schrieffer and F. Wilczek, Phys. Rev. Lett. **53**, 722 (1984), F. Wilczek and A. Zee, Phys. Lett. **147 B**, 325 (1984), F. D. M. Haldane and Y. S. Wu, Phys. Rev. Lett. **55**, 2887 (1985), R. B. Laughlin, Science **242**, 525 (1988), Ann. Phys. (N. Y.), **191**, 163 (1989), X. G. Wen, F. Wilczek and A. Zee, Phys. Rev. **B 39**, 11413 (1989).

[2] Y. S. Wu, Phys. Rev. Lett. **52**, 2103 (1984).

[3] Y. S. Wu, Phys. Rev. Lett. **53**, 111 (1984).

[4] Y. H. Chen, F. Wilczek, E. Witten and B. I. Halperin, Int. J. Mod. Phys. **B 3**, 1001 (1989).

[5] R. Jackiw, Ann. Phys. (N. Y.), **201**, 83 (1990).

[6] J. M. Leinaas and J. Myrheim, Nuovo Cimento, **37 B**, 1 (1977).

[7] C. Chou, Phys. Rev. **D 44**, 2533 (1991).

[8] A. A. Chin and C. R. Hu, Phys. Rev. Lett. **69**, 229 (1992).

[9] M. D. Johnson and G. S. Canright, Phys. Rev. **B 41**, 6870 (1990), G. V. Dunne, A. L. Lerda and C. A. Trugenberger, *Mod. Phys. Lett. A 6*, 2819 (1991), K. H. Cho, C. Rim and D. S. Soh, Phys. Lett. **A 164**, 65 (1992), A. Dasnieres de Veigy and S. Ouvry, Phys. Rev. Lett. **72**, 600 (1994).

[10] G. S. Canright and M. D. Johnson, J. Phys. **A**: Math. Gen., **27**, 3579; S. He, X. C. Xie and F. C. Zhang, Phys. Rev. Lett. **68**, 3460 (1992).

[11] Ian I. Kogan, Int. J. Mod. Phys. **A9**, 3887 (1994).

[12] P. B. Wiegmann and A. V. Zabrodin, cond-mat/9310017; cond-mat/9312088.

[13] C. N. Yang and Mo-lin Ge, *Quantum Group*, Nankai University Proceedings, references therein.

[14] L. D. Faddeev and R. M. Kashaev, hep-th/9312133.

[15] E. K. Sklyanin, Func. Anal. App. **17**, 273 (1983).

[16] J. Zak, Phys. Rev. **134**, 6A (1964).

[17] L. D. Landau, Z. Phys. **64**, 629 (1930).